X100/301

NATIONAL QUALIFICATIONS 2007 TUESDAY, 15 MAY 9.00 AM - 10.10 AM

MATHEMATICS HIGHER Units 1, 2 and 3 Paper 1

(Non-calculator)

Read Carefully

- 1 Calculators may NOT be used in this paper.
- 2 Full credit will be given only where the solution contains appropriate working.
- 3 Answers obtained by readings from scale drawings will not receive any credit.





FORMULAE LIST

Circle:

The equation $x^2 + y^2 + 2gx + 2fy + c = 0$ represents a circle centre (-g, -f) and radius $\sqrt{g^2 + f^2 - c}$. The equation $(x - a)^2 + (y - b)^2 = r^2$ represents a circle centre (a, b) and radius r.

Scalar Product:

 $a.b = |a| |b| \cos \theta$, where θ is the angle between a and b

or
$$\mathbf{a}.\mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$$
 where $\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$.

Trigonometric formulae:

$$\sin (A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos (A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\sin 2A = 2\sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$= 2\cos^2 A - 1$$

$$= 1 - 2\sin^2 A$$

Table of standard derivatives:

f(x)	f'(x)
sin ax	a cos ax
cos ax	$-a\sin ax$

Table of standard integrals:

$$f(x) \qquad \int f(x) dx$$

$$\sin ax \qquad -\frac{1}{a}\cos ax + C$$


$$\cos ax \qquad \frac{1}{a}\sin ax + C$$

ALL questions should be attempted.

Marks

3

1. Find the equation of the line through the point (-1, 4) which is parallel to the line with equation 3x - y + 2 = 0.



2. Relative to a suitable coordinate system A and B are the points (-2, 1, -1) and (1, 3, 2) respectively.

A, B and C are collinear points and C is positioned such that BC = 2AB.

Find the coordinates of C.

3. Functions f and g, defined on suitable domains, are given by $f(x) = x^2 + 1$ and g(x) = 1 - 2x.

Find:

- (a) g(f(x));

- 2
- (b) g(g(x)). 2


4. Find the range of values of k such that the equation $kx^2 - x - 1 = 0$ has no real roots.

5. The large circle has equation $x^2 + y^2 - 14x - 16y + 77 = 0.$

> Three congruent circles with centres A, B and C are drawn inside the large circle with the centres lying on a line parallel to the x-axis.

> This pattern is continued, as shown in the diagram.

> Find the equation of the circle with centre D.



5

[Turn over

6. Solve the equation $\sin 2x^{\circ} = 6\cos x^{\circ}$ for $0 \le x \le 360$.

4

7. A sequence is defined by the recurrence relation


$$u_{n+1} = \frac{1}{4}u_n + 16, \ u_0 = 0.$$

(a) Calculate the values of u_1 , u_2 and u_3 .

3

Four terms of this sequence, u_1 , u_2 , u_3 and u_4 are plotted as shown in the graph.

As $n \to \infty$, the points on the graph approach the line $u_n = k$, where k is the limit of this sequence.



2 3 4 *n*

(ii) Find the exact value of k.

3

- 8. The diagram shows a sketch of the graph of $y = x^3 4x^2 + x + 6$.
 - (a) Show that the graph cuts the x-axis at (3, 0).
 - (b) Hence or otherwise find the coordinates of A.



(c) Find the shaded area.

5

1

- **9.** A function f is defined by the formula $f(x) = 3x x^3$.
 - (a) Find the exact values where the graph of y = f(x) meets the x- and y-axes.
 - (b) Find the coordinates of the stationary points of the function and determine their nature.
 - (c) Sketch the graph of y = f(x).

 u_n

k

7

2

Marks

10. Given that $y = \sqrt{3x^2 + 2}$, find $\frac{dy}{dx}$.

- 3
- 11. (a) Express $f(x) = \sqrt{3}\cos x + \sin x$ in the form $k\cos(x a)$, where k > 0 and $0 < a < \frac{\pi}{2}$.
- 4

4

(b) Hence or otherwise sketch the graph of y = f(x) in the interval $0 \le x \le 2\pi$.

[END OF QUESTION PAPER]

X100/303

NATIONAL QUALIFICATIONS 2007 TUESDAY, 15 MAY 10.30 AM - 12.00 NOON MATHEMATICS HIGHER Units 1, 2 and 3 Paper 2

Read Carefully

- 1 Calculators may be used in this paper.
- 2 Full credit will be given only where the solution contains appropriate working.
- 3 Answers obtained by readings from scale drawings will not receive any credit.





FORMULAE LIST

Circle:

The equation $x^2 + y^2 + 2gx + 2fy + c = 0$ represents a circle centre (-g, -f) and radius $\sqrt{g^2 + f^2 - c}$. The equation $(x - a)^2 + (y - b)^2 = r^2$ represents a circle centre (a, b) and radius r.

Scalar Product:

 $a.b = |a| |b| \cos \theta$, where θ is the angle between a and b

or
$$\boldsymbol{a}.\boldsymbol{b} = a_1b_1 + a_2b_2 + a_3b_3$$
 where $\boldsymbol{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$ and $\boldsymbol{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$.

Trigonometric formulae:

$$\sin (A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos (A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\sin 2A = 2\sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$= 2\cos^2 A - 1$$

$$= 1 - 2\sin^2 A$$

Table of standard derivatives:

f(x)	f'(x)
sin ax	a cos ax
cos ax	$-a\sin ax$

Table of standard integrals:

$$f(x) \qquad \int f(x) dx$$


$$\sin ax \qquad -\frac{1}{a}\cos ax + C$$

$$\cos ax \qquad \frac{1}{a}\sin ax + C$$

1. OABCDEFG is a cube with side 2 units, as shown in the diagram.

B has coordinates (2, 2, 0).

P is the centre of face OCGD and Q is the centre of face CBFG.



(a) Write down the coordinates of G.

1


(b) Find p and q, the position vectors of points P and Q.

2

(c) Find the size of angle POQ.

5

2. The diagram shows two right-angled triangles with angles c and d marked as shown.



(a) Find the exact value of $\sin(c+d)$.

4


- (b) (i) Find the exact value of $\sin 2c$.
 - (ii) Show that $\cos 2d$ has the same exact value.

4

3. Show that the line with equation y = 6 - 2x is a tangent to the circle with equation $x^2 + y^2 + 6x - 4y - 7 = 0$ and find the coordinates of the point of contact of the tangent and the circle.

6

- 4. The diagram shows part of the graph of a function whose equation is of the form $y = a\sin(bx^{\circ}) + c$.
 - (a) Write down the values of a, b and c.
 - (b) Determine the exact value of the x-coordinate of P, the point where the graph intersects the x-axis as shown in the diagram.




3

3

[Turn over

5. A circle centre C is situated so that it touches the parabola with equation $y = \frac{1}{2}x^2 - 8x + 34$ at P and Q.



(a) The gradient of the tangent to the parabola at Q is 4. Find the coordinates of Q.

5

(b) Find the coordinates of P.

2

(c) Find the coordinates of C, the centre of the circle.

2

6. A householder has a garden in the shape of a right-angled isosceles triangle.

10 m 10 m Decking

Side Wall

It is intended to put down a section of rectangular wooden decking at the side of the house, as shown in the diagram.

- (i) Find the exact value of ST.
- - (ii) Given that the breadth of the decking is x metres, show that the area of the decking, A square metres, is given by

$$A = \left(10\sqrt{2}\right)x - 2x^2.$$

(b) Find the dimensions of the decking which maximises its area.

5

7. Find the value of $\int_0^2 \sin(4x+1) dx$.

4

The curve with equation $y = \log_3(x - 1) - 2.2$, where x > 1, cuts the x-axis at the point (a, 0).

Find the value of a.

4

2


2

9. The diagram shows the graph of $y = a^x$, a > 1.

On separate diagrams, sketch the graphs of:




(b)
$$y = a^{1-x}$$
.



The diagram shows the graphs of a cubic function y = f(x) and its derived function y = f'(x).

Both graphs pass through the point (0, 6).

The graph of y = f'(x) also passes through the points (2, 0) and (4, 0).



- (a) Given that f'(x) is of the form k(x-a)(x-b):
 - (i) write down the values of a and b;
 - (ii) find the value of k.

- 3
- (b) Find the equation of the graph of the cubic function y = f(x).

- Two variables x and y satisfy the equation $y = 3 \times 4^x$. 11.
 - (a) Find the value of a if (a, 6) lies on the graph with equation $y = 3 \times 4^x$.
- 1

(b) If $(-\frac{1}{2}, b)$ also lies on the graph, find b.

- 1
- (c) A graph is drawn of $\log_{10}y$ against x. Show that its equation will be of the form $\log_{10} y = Px + Q$ and state the gradient of this line.

[END OF QUESTION PAPER]